

THE UNKLED SHAWES OF AMERICA

TO ALL TO WHOM THESE: PRESENTS: SHALL COME;

Seed Research of Pregon

MICEONS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HERS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE CHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR USING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE REPOSES, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

FESCUE, TALL

'Rendition'

In Cestimonn Marrest, I have hereunto set my hand and caused the seal of the Hant Haristy Frotestion Office to be affixed at the City of Washington, D.C. this twelfth day of September, in the year two thousand and seven.

Allost:

acrose

Commissioner Plant Variety Protection Office Agricultural Marketing Service Sucretary of

REPRODUCE LOCALLY, include form number and date on all re-	roductions	Form Approved - CMB No. 0581-0055			
U.B. DEPARTMENT OF AGRICULTURAL MARKETING SCIENCE AND TECHNOLOGY - PLANT VARIE APPLICATION FOR PLANT VARIETY PROT (Instructions and Information collection burde	SERVICE TY PROTECTION OFFICE ECTION CERTIFICATE	The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552e) and the Paperwork Reduction Act (PPA) of 1995. Application is required in order to determine if a plant variety protection cartificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426).			
1. NAME OF OWNER	, tracounters ou savetee)	2. YEMPORARY DESIGNATION OR	3. VARIETY NAME		
Seed Research of Ore	gon	SRX 8V9	Rendition		
4. ADDRESS (Street and No., or R.F.D. No., City, State, and Z	P Code, and Country)	5. TELEPHONE (Include area code)	FOR OFFICIAL USE ONLY		
27630 Llewellyn Rd.	•	541-757-2663	PYPO NUMBER		
Corvallis, OR 97333		6. FAX (include sine code) 541-752-2065	200300174 FLINGDATE		
7. If the Owner Name is NOT A Person, Give Form ORGANIZATION (corporation, perhership, escoclation, etc.) Corporation	O R	8. DATE OF INCORPORATION 1983	2/26/2003		
10. NAME AND ADDRESS OF OWNER REPRESENTATIVES Dr. Leah A. Brilman	TO SERVE IN THIS APPLICATION. (First	person listed will receive all papers)	F FRING AND EXAMINATION FEES:		
Seed Research of Orego	n		19 6 . 1		
27630 Llewellyn Rd.	•	•	R DATE 2/26/2003		
•			e . 768,00		
Corvallis, OR 97333			E DATE 8/28/2007		
11. TELEPHONE (Include area code)	12. FAX (include area code)	13. E-MAIL	14. CROP KIND (Common Name)		
541-758-9115	541-752-2065	srofarm@attgloba	l.net Tall fescue		
15. GENUS AND SPECIES NAME OF CROP		18. FAMILY NAME (Botanical)	17. IS THE VARIETY A FIRST GENERATION		
Festuca arundinacea		Poageae	HYBRID?		
18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT (Follow instructions on reverse)	SUBMITTED	19. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? See Section 53(e) of the Plant Variety Protection Act)			
a. DE Exhibit A. Origin and Breeding History of the Varie	v	□ YES (If "yes", answer items 20 and 21 below) 原 NO (If "no", go to item 22)			
b. K Eichibk B. Statement of Distinciness		20. DOES THE OWNER SPECIFY THAT SEED OF THIS I YES IND VARIETY BE LIMITED AS TO NUMBER OF CLASSES?			
c. PL Exhibit C. Objective Description of Variety			_		
d. Schlöt D. Additional Description of the Variety (O		IF YES, WHICH CLASSES? D FOUNDATION D REGISTERED C CERTIFIED			
Bi Exhibit E. Statement of the Beats of the Owner's C Bi Voucher Sample (2,500 visible uninvaled seeds or verification that tissue culture will be deposited an	for tuber propagated varieties,	21. DOES THE OWNER SPECIFY THAT SEED OF THIS TYPES NO VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? IF YES, SPECIFY THE NUMBER 1,2,3,46., FOR EACH CLASS.			
g. 15 Filing and Examination Fee (\$3,652), made payab States" (Net to the Plant Variety Protection Office)	e to "Tressurer of the United	☐ FOURDATION ☐ RESISTERED ☐ CERTIFIED (If additional explanation is necessary, please use the space indicated on the reverse.)			
22. HAS THE VARIETY (INCLUDING ANY HARVESTED MAT FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRAI OR		23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)?			
OTHER COUNTRIES? YES IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALUSE BOR EACH COUNTRY AND THE DROUMSTANCES. (P		[] YES \$\frac{12}{12}\$ NO IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Please use space indicated on reverse.)			
24. The owners declare that a visible sample of basic seed of	······································		and a not with a sub-property of the second		
for a fother propagated variety a finance culture will be dep	celled in a public repository and entirtainer	d for the duration of the certificate.			
The undersigned owner(s) is(are) the owner of this secur and is entitled to protection under the provisions of Sections	on 42 of the Plant Variety Profestion Act.		stinct, uniform, and stable as required in Section 42,		
Owner(s) le(sre) informed that take representation hersi	r can propercize protection and least in per				
Plan a. Bulm	an	SIGNATURE OF OWNER			
NAME (Pitranea puto) or type)		NAME (Please print of type)			
Leah A. Brilman	1				
CAPACITY OR TITLE	DATE	CAPACITY OR TITLE	DATE		
Research Director	2-25-03		· -		

\$2-470 (02-10-2003) designed by the Plant Variety Projection Office using Word 2000. Replaces former various of \$1-470, which are ob-

INSTRUCTIONS

2 0 0 7 0 0 1 2 6

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPO: (1) Completed application form signed by the owner, (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$3,652 (\$432 filing fee and \$3,220 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfiled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$432 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

> **Plant Variety Protection Office** Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM

- 18a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
 - (2) the details of subsequent stages of selection and multiplication;
 - (3) evidence of uniformity and stability; and
 - (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified.)

22. CONTINUED FROM FRONT (Please provide the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.)

August 30, 2002

23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the variety is protected by intellectual property right (Plant Breeder's Right or Patent).)

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act. and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center--East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to averaging the collection of information, instructions, searching existing data sources, gethering and maintaining the data needed, and completing and reviewing the collection of information. ted to average 3.0 hours per response, including the time for reviewing

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, gender, religion, age, disability, sexual orientation, marital or family status, political beliefs, parental status, or protected genetic information. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotage, etc.) should contact USDA's TARGET Center at 202-720-2600 (voice and TDD).

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or cell 202-720-5964 (voice and USDA is an equal opportunity provider and employer.

ST-470 (02-10-2003) design with Word 2000. Replaces former versions of ST-470, which are obsolete

Exhibit A.

Origin and Breeding History of Rendition (SRX 8V9) Tall Fescue

Rendition (SRX 8V9) tall fescue (Festuca arundinacea Schreb.) is a low-growing, dark green, fine-leaved, turf-type tall fescue selected from the maternal progenies of 4 different clones.

The parental germplasm of Rendition tall fescue traces its origin to plants selected from old turfs of the United States in a germplasm collection program initiated in 1962 and to plants selected from or related to Rebel tall fescue (Funk et al., 1981). Attractive clones were selected from old turfs in Birmingham, AL; Athens, Atlanta, and Millegeville, GA; Preston, ID; Baltimore, MD; Bayonne, Jersey City, Elizabeth, Princeton, and Cape May, NJ; eastern North Carolina; Philadelphia, PA; Nashville, TN; Lexington, KY; Cincinnati, OH; Dallas, TX; and northern Mississippi. The tall fescue plants selected from old turfs were of unknown origin. All were large patches of turf surviving in stressful environments indicating that they had persisted and developed over a period of many years.

A few hundred attractive, turf-type plants were collected and established in spaced-plant nurseries and/or frequently mowed clonal evaluation trials at Rutgers University. All but a few dozen of the most promising plants were quickly discarded. The best selections were very different from any tall fescue variety in existence at the time of collection. They produced lower-growing turfs with finer leaves, greater density, darker color, and greater tolerance of close mowing.

The most promising plants were identified by their persistence and appearance in old turfs and their performance in spaced-plant nurseries, mowed clonal evaluation tests, and single-plant progeny trails under turf maintenance. Intercrosses of the best performing plants were subjected to varying cycles of phenotypic and genotypic

selection depending on their date of collection. New sources of germplasm were added to the breeding program as it became available from the continuing collection program. Each cycle of selection showed continued progress in producing lower-growing, darker green, attractive plants with improved turf performance scores. Selection was also effective in maintaining high seed yields, and good stress tolerance. Substantial progress was made in developing tall fescues with finer leaves, a lower growth profile, increased persistence under close mowing, and increased density.

Large numbers of single-plant progenies were seeded in turf evaluation trials at the Plant Science Research Farm at Adelphia, NJ in 1995. The plants selected for progeny evaluation were selected from spaced-plant nurseries at Adelphia following varying cycles of phenotypic and genotypic selection of germplasm selected from old turfs and germplasm selected from or related to Rebel tall fescue.

Following a period of summer stress due to heat, drought, and disease, a total of 1,900 plants were selected from 19 of the best performing single-plant progeny turf plots. All 19 progenies were from the 1995 test. Selection of progenies was based on performance records, as well as appearance at the time the plants were selected from these progeny plots. Selection of plants from each progeny was based on an attractive dark green color, fine leaves, abundant tillering, and freedom from disease. Selected plants were sent to Seed Research of Oregon in the fall of 1996. For most lines received 70 plants per line were established, with the exception of A95-246 where 105 plants were established and only 35 of A95-212 and A95-262. In 1997 and 1998, Seed Research of Oregon classified the plants received from Rutgers based on maturity, height, color and leaf texture. The plants designated at double dwarf and medium were all moved to a separate block in 1996 and allowed to interpollinate. This included progeny from A95-514, A95-246, A95-539 and A95-502. The resulting progeny for each clone was sent to

Rutgers for evaluation, plus progeny of two plants from A95-539, designated W77-17E and W77-24E were also sent. This block was rouged for seed yield, uniformity of maturity and stem rust before pollination. These plants were all very dark green with fine leaf texture. A total of 21 plants from A95-514, 24 plants from A95-539, 18 from A95-502 and 23 from A95-246 were harvested. In 1997 a new breeder block was established of 105 progeny from each of the following lines, W77-17E (A95-539), W17-24E(A-95-539), W78 (A95-246) and W59 (A95-514). The performance of A95-502 caused it to be eliminated. After rouging for uniformity the following progeny remained of each line, W59, 63 plants, W78, 53 plants, W77-17E, 82 plants and W77-24E, 72 plants. All of these lines had high endophyte levels. These were harvested for breeder seed in 1999.

The first foundation field was established in the fall of 2000. A few variants that are taller and with wider leaves than the rest of the variety have been observed in foundation field (less than 1%), but have never been observed in our PVP nurseries of seventy plants. The first foundation field was established in the fall of 1999 along with turf plots of the final composite and the initial PVP nursery. The same percentage of variants have been found in certified production, although the taller plants were rouged from the foundation field. Three generations of increase are approved. Rendition has been shown to be uniform and stable in foundation and certified production.

References

- Buckner, R. C., J. B. Powell, and R. V. Frakes. 1979. Historical Development, in Buckner, R. C., and L. P. Bush (editors) Tall Fescue. Agronomy Monograph 20. American Society of Agronomy, Crop Science Society of America, Soil Science Society of America, Inc., Publishers. Madison, Wisconsin pages 1-8.
- 2. Funk, C.R., R.E. Engel, W.K. Dickson, and R.H. Hurley. 1981. Registration of Rebel tall fescue. Crop Sci. 21:632.

200300174

Diagram of Origin and Breeding History of Rendition Tall Fescue

1. 1962 to 1994

Germplasm collection, evaluation, and genetic improvement.

2. 1991 to 1995

Planted single-plant progenies of plants selected from current cycles of population improvement programs in closely mowed turf trials at Adelphia and North Brunswick, NJ.

3. 1996

Selected 1,900 plants from 19 of the best performing single-plant progeny turf plots planted in 1995. Selected plants were sent to Seed Research of Oregon for further selection and classification. Plants were selected and moved according to growth habit, fine leaf texture, freedom from disease and seed yield.

4. 1997

New breeder block established from progeny of 3 clones (only from two plants of one clone). Breeder block rouged for uniformity, freedom from disease and seed yield.

Each plant of Rendition tall fescue traces at least 20 percent of its ancestral germplasm to plants selected from or related to Rebel tall fescue.

200300174

EXHIBIT B.

'RENDITION' TALL FESCUE NOVELTY STATEMENT

'Rendition' tall fescue most closely resembles SR 8600 but can be distinguished from this variety by a combination of the following:

- 1) Rendition has a significantly greater plant width than SR 8600 in 2000 and 2001 (Tables 3 and 4).
- 2) Rendition has a significantly lesser panicle length than SR 8600 in 2000 and 2001 (Tables 3 and 4).

U.S. DEPARTMENT OF AGRICULTURE PLANT VARIETY PROTECTION OFFICE, AMS, USDA NATIONAL AGRICULTURAL LIBRARY Bldg., Rm. 500 10301 BALTIMORE Blvd. BELTSVILLE, MD 20705

OBJECTIVE DESCRIPTION OF VARIETY TALL & MEADOW FESCUES

(Festuca spp.)

NAME OF APPLICANT(S) Seed Research of Ore	_	ORARY DESIGN. (8V9		LIETY NAME ndition
ADDRESS (Street and No., or R.F.D. No., City, State, an 27630 Liewellyn Rd.	d ZIP Code)		· ·	OFFICIAL USE ONLY O NUMBER
Corvallis, OR 97333			200	500174
Place the appropriate number that describes the varietal char 089). Characteristics described, including numerical meass should be for SPACED PLANTS. Royal Horticultural Characteristics marked with an asterisk * are characteristics	urements, should re Society or any re	epresent those that cognized color far	are typical fo	r the variety. Measured data
* 1. SPECIES: (With comparison varieties, use varieties wi	ithin the species of	the application varie	ety)	
$X_1 = F$. arundinacea (Tall)	Turf Types			
	olympic 4 = Bor ebel Jr. 10 = M		5 = Arid 11 = Crewcut	6 = Rebel II 12 = Bonsai
	Forage Types			
	Martin AU Triumph		23 = Mozark 27 = Cajun	
2 = F. pratensis (Meadow)				
30 = Admira 31 = Beaumon	t 32 = Comtessa	33 = Ensign	34 = Trader	
* 2. CYTOLOGY:				
42 Chromosome Numb	ег			
3. ADAPTATION: (0 = Not Tested; 1 = Not Adapted; 2 =	Adapted)			
2 Transition Zone 2 West 2	Northeast(Other (Specify):		
* 4. MATURITY: (Date First Headed, 10% of Panicle Eme	ergence)			
	U Triumph ate (Silverado)		4 = K31, Kenl 9 = Very late	ny 5 = Medium (Rebel)
Date Headed 113.9 2001 data Locati	ion <u>Corvallis. C</u>	R	,	_
Maturity same as4_ Comparison Va	ariety			
Days later than				

* 5. MATURE PLANT HEIGHT from crown to top of panicle, it		* INTERNODE LENGT (First internode subtendin	H CM () () () () () () () () () (
<u>95.0</u> cm Height		42.1 cm Internode le	ngth
22.0 cm shorter than	9	6.9 cm shorter than	_Titan_
Height same as	_8_ Comparison Variety	Length same as	Comparison variety
cm taller than	_	cm longer than	
* HEIGHT AT EAR EMERGE	ENCE CM: (Flag leaf height from c	rown to flag leaf node)	
47.1 cm Height			
5.8 cm shorter tha	n <u>Z</u>		
Height same as	Comparison Variety		
cm taller than			
* 6. GROWTH HABIT: (Mature	Plants)		
9 1 = Prostrate (7 = Semierect (•	•	rizontal ()
* 7. RHIZOMES (Psuedo):		***************************************	
mm Length	3 1 = Absent () $2 = Ra$	are (Rebel) 3 = Cor	mmon ()
* 8. LEAF BLADE: (Tiller leaves	s/ turf color)		
7 = Me	ght green () $3 = M$ edium dark green () $9 = Ve$ y rating of comparison variety SR 8	ery dark green ()	
* 9 Anthocyanin:			·
*_1 Basal Hairs: 1 = Ab	sent () 9 = Present ()	
*4.8 Margins:	1 = Smooth ()	5 = Semi-rough ()	9 = Rough ()
*_8_Width Class:	1 = Very coarse () 3 = Co 7 = Fine () 9 = Ve		dium ()
* TILLER LEAF LENGTH CM:	(First leaf subtending the flag leaf)	* TILLER LEAI	F WIDTH MM:
26.5 cm Tiller Leaf	Length	_6.4_ mm Tille	er Leaf Width
10.2 cm shorter than	10	1.4 mm narro	ower than <u>Titan</u>
Length same as	Comparison Variety	Width same as	Comparison variety
cm longer than		mm wider	than

8. LEAF BLADE: (continued)		GAG GAG GAG	
FLAG LEAF LENGTH CM:		FLAG LEAF WIDTH MM. C S C C S 7 4	
14.0 cm Flag Lea	f Length	3.9 mm Flag Leaf Width	
cm shorter than		1.7 mm narrower than 9	
Length same as	Titan Ltd. Comparison Variety	Width same as <u>Titan Ltd.</u> Comparison var	iety
cm longer than		mm wider than	
* 9. LEAF SHEATH: (Basal Po	ortion)		
*_9 Anthocyanin (seed	dling): $1 = \text{Absent (K31)}$	9 = Present ()	
*4.2 Auricle Hairiness	s: $I = Absent ()$	9 = Present ()	
* 10. PANICLE: (At seed matur	rity except where noted.)		
*4.0 Shape: $1 = N$	Narrow-tapering () $5 = 0$	vate () $7 = Oblong () 9 = Other (specify)$	
*_ 4.0 Type: 1 = 0	Compact (appressed) 5 = In	termediate () 7 = Open () 9 = Other (specify)	
*_8_ Orientation:	1 = Nodding ()	9 = Erect ()	
*_9 Branch Pubescence	æ: 1 = Glabrous (32%)	9 = Pubescent (68%)	
* 1 Anther Color (At	t anthesis): 1 = Yellowish Green 4 = Purplish	2 = Green 3 = Bluish Green (68% 1, 32% 4) 5 = Reddish 6= Other (Specify)	
*_2_ Glume Color (At	t anthesis): 1 = Yellowish Green 4 = Purplish	2 = Green 3 = Bluish Green (84% 2, 16% 5) 5 = Reddish 6= Other (Specify)	
*18.3 cm Panicle Len	igth (from base to tip, if nodding, stra	ighten; after anthesis)	
5.5 cm shorter than	SR 8600		
Length same as	12 Comparison Variety		
cm longer than			
* 11. SEED: (With Lemma & P	Pelea)		
* <u>2639</u> mg per 1000 s	eeds		
mg less than			
Weight same as	SR 8600 Comparison Variety		
447 mg more than	<u>Titan</u>		
PALEA: (Keels or Margins)	4.5 Hairs: 1 = Absent () 5 = Short (Missouri 96) 9 = Long ()	
LEMMA:	2.7 Hairs: 1 = Absent (Ke	nhy) 5 = Several () 9 = Many (Missour	ri 96)
6.1 mm Lemma Lengtl	h (Mature)	1.2 mm Lemma width	
0.3 mm shorter than	<u>\$R 8600</u>	0.2 mm narrower than SR 8600	
Length same as	Comparison Variety	Width same as Comparison variety	
0.3 mm longer than	Titan Ltd.	0.2 mm wider than Titan	

10. PANICLE: (continued)			200300176	
*AWNS:	9 AWNS:	l = Absent () 9 = Present (Falcon)	· · · · · · · · · · · · · · · · · · ·	
1.1 mm Awn len	ngth (Of those pre	esent.)			
mm Shorte	er than				
Length same as	SR 86	600 Comparisor	Variety		
mm Longer	r than				
12. DISEASE, INSECT, A	AND NEMATOR	DE REACTION	(0= Not Tested 1= Lo	east Resistant 9= Most Resistant)	_
Melting-out I	Drechslera poae		Blind S	Seed Gloeotinia temulenta	
<u>5.7</u> Leaf Spot D.	siccans		Dollar	Spot Lanzia, Mollerdiscus spp.	
Net Blotch D). dictyoides		Stem F	Rust Puccinia graminis	
5.8 Brown Patch	n Rhizoctonia sold	ani	T. Blig	ght <i>Typhula incarnata</i>	
C. Leaf Spot	Cercospora fectu	сае	Pythiu	ım Blight <i>Pythium</i> spp.	
Pink Snow M	Aold <i>Gerlachia ni</i>	ivalis	Powde	ery Mildew <i>Erysiphe graminis</i>	
Silver Top F.	ξ trivinctum, F . r	oseum	Crown	n Rust Puccinia coronala	
Other Disease	e			· · · · · · · · · · · · · · · · · · ·	
Other Insect _					
Other Nemato	ode				

14. GIVE VARIETY OR VARIETIES THAT MOST CLOSELY RESEMBLE THE APPLICATION VARIETY. For the following characteristics, indicate the degree of resemblance with the following scale:

5 = Tolerant (

5 = Tolerant (

5 = Tolerant (

9 = Resistant (

9 = Resistant (

9 = Resistant (

1 = Application variety is less than comparison variety 2 = Same as 3 = More than, better, greater, darker, etc.

1 = Susceptible (

1 = Susceptible (

1 = Susceptible (

13. ENVIRONMENTAL STRESS

___ Drought Stress

Shade Stress

Winter Stress

Character	Varieties	Rating	Character	Varictics	Rating	
Leaf Width	SR 8600	1	Leaf Color	SR 8600	1	
Paniele Color	SR 8600 (Red)	3	Paniele Shape	SR 8600 (Na	rrow) 3	
Seed Size	SR 8600	1	Cold Injury			
Winter Color			Heat			
Disease			Plant Width	SR 8600	3	

^{* 15.} EXPERIMENTAL: Give a brief summary of the experimental design utilized to collect the data used on this form. Cultural conditions, number of plants measured and plant spacing must be specified. Spaced plants, 3 ft. spacing. 2 replications, 35 plants each, randomized complete block, Tilt for stem rust, Diuron + Goal weed control

EXHIBIT D. ADDITIONAL DESCRIPTION OF THE VARIETY




Table 1. Heading and anthesis dates for tall fescue varieties in 2000. The plants were established in the fall of 1999 with the seed of control varieties obtained from the repository at the Plant Material Center at Pullman, WA or from the breeder. Two replications of 35 plants were established and the number of plants with at least three heads are as shown. Two applications of TILT were applied to the PVP block to control stem rust. Means followed by the same letter are not significantly different at the 0.05 level.

Variety	Plant Number			Anthesis <u>Mear</u>	
KY-31 NE	68	110.0 a	ı	145.2 ab)
KY-31 E	68	110.9 a	l	146.3 k	oc
Rendition	70	111.4 a	ı	144.2 a	
Titan	69	114.2	b	147.6	С
SR 8600	69	116.1	bc	149.7	е
Rebel li	69	116.8	С	148.9	de
MiniMustang	68	117.3	cd	151.1	f
Crewcut	67	119.5	de	152.0	fg
Silverado	69	120.0	ef	152.9	g
Grande	68	120.5	ef	152.3	fg
SR 8500	70	120.6	ef	151.7	fg
Rebel Jr.	69	120.6	ef	152.5	fg
Shortstop	67	121.5	efg	153.1	gh
Bonanza	62	121.7	efg	152.7	g
SR 8210	67	121.7	efg	152.2	fg
Crewcut II	67	121.8	fg	152.6	g
SR 8200	70	123.0	g	152.9	gh
Bonsai	66	123.3	g	154.3	h
LSD @5%		2.3		1.4	




Table 2. Heading and anthesis dates for tall fescue varieties in 2001. The plants were established in the fall of 1999 with the seed of control varieties obtained from the repository at the Plant Material Center at Pullman, WA or from the breeder. Two replications of 35 plants were established and the number of plants with at least three heads are as shown. These are the same plants used for data collection in 2000. Two applications of TILT were applied to the PVP block to control stem rust. Means followed by the same letter are not significantly different at the 0.05 level.

Variety	Plant Number	Heading Date <u>Mean</u>	Anthesis Date Mean
KY-31 NE	64	111.6 a	148.8 ab
KY-31 E	6 9	113.3 ab	149.6 bc
Titan Ltd.	70	113.9 b	147.3 a
Titan	66	116.4 c	149.1 bc
Rebel li	69	116.9 c	150.2 bcd
SR 8600	66	117.1 c	148.9 abc
Rendition	69	117.8 c	150.5 cde
SR 8210	67	120.2 d	152.1 ef
Silverado	69	120.3 d	151.9 def
SR 8500	68	120.4 d	151.7 fg
Crewcut II	66	120.7 de	153.0 fgh
Shortstop	69	121.1 def	153.9 ghij
MiniMustang	65	121.3 def	153.1 fgh
Crewcut	66	122.5 efg	152.5 fg
SR 8200	64	122.7 fg	154.7 hij
Rebel Jr.	69	123.5 gh	155.5 j
Bonanza	62	123.6 gh	152.9 fg
Bonsai	63	123.8 gh	153.4 fghi
Grande	68	124.9 h	155.1 ij
LSD @5%		1.9	1.7

Table 3. Comparative plant morphological measurements of tall fescue varieties in a spaced plant nursery in 2000 near Corvallis, OR. The plants are those described in Table 1. Randomly selected reproductive tillers were selected after anthesis for measurement, with 60 measurements for each characteristic, 30 per rep. Panicle length in this table is that measured in the field on the same tillers used for plant height.

Variety	Plant Height (cm)	Panicle Branch Internode Length (cm)	Height at Ear Emergence (cm)	Plant Width (cm)	Panicle Length (cm)
Bonsai	104.30	23.52	57.83	31.35	20.58
Crewcut II	106.82	21.68	62.43	29.29	23.50
Silverado	109.72	23.45	63.78	31.70	24.57
SR 8500	112.42	22.74	65.64	29.90	24.57 25.23
Rendition	114.40	25.97	63.95	34.77	25.23 22.47
MiniMustang	115.34	23.91	65.85	35.23	25.85
SR 8600	118.11	25.01	63.80	29.13	26.34
Grande	120.34	26.24	69.77	35.27	28.09
Rebel Jr.	122.22	26.34	70.03	31.68	28.22
Crewcut	122.59	25.63	71.00	36.38	27.75
Shortstop	122.63	24.45	71.36	33.83	26.98
Titan Ltd.	123.22	24.82	72.66	32.74	24.76
SR 8200	126.18	25.97	73.81	33.23	28.91
SR 8210	128.59	27.34	78.06	32.65	28.15
Bonanza	132.91	28.35	79.18	34.59	32.89
Rebel II	135.90	27.95	82.05	37.58	31.24
Titan	136.86	26.97	81.91	32.35	30.81
Kentucky 31 NE	149.21	28.79	90.13	40.08	33.37
Kentucky 31 E	149.63	27.82	92.40	36.98	35.05
LSD @5%	5.11	1.73	4.48	2. 4 5	2.02

Height at ear emergence = flag leaf height

200000174

Table 4. Comparative plant morphological measurements of tall fescue varieties in a spaced plant nursery in 2001 near Corvallis, OR. The plants are those described in Table 2. Randomly selected reproductive tillers were selected after anthesis for measurement, with 60 measurements for each characteristic, 30 per rep. Panicle length in this table is that measured in the field on the same tillers used for plant height. These are the same plants used in 2000 but not as many experimental varieties were measured in 2001.

Variety	Plant Height (cm)	Panicle Branch Internode Length (cm)	Height at Ear Emergence (cm)	Plant Width (cm)	Panicle Length (cm)
Rendition	95.00	42.10	47.14	48.51	18.29
Silverado	104.33	40.63	52.86	44.49	20.80
Bonsai	106.74	41.87	56.22	39.44	18.04
SR 8600	108.41	39.75	54.54	44.06	23.87
Titan Ltd.	112.59	44.03	62.78	48.19	22.90
MiniMustang	116.29	45.73	56.70	40.86	22.59
Rebel Jr.	117.09	44.00	56.59	42.05	21.69
Shortstop	117.68	43.32	52.86	44.37	26.02
Titan	122.88	48.96	69.39	43.43	24.44
Crewcut	124.35	45.92	63.87	45.01	25.63
Kentucky 31 E	144.47	56.06	76.94	43.62	27.10
LSD @5%	5.15	4.33	5.27	3.40	3.40

Height at ear emergence = flag leaf height

Table 5. Leaf characteristics of tall fescue varieties in 2000 near Corvallis, OR. The plants are those described in Tables 1 and 3. The leaves were measured on the reproductive tillers utilized in Table 3. The vegetative leaf was measured as the last fully expanded leaf on vegetative tillers on the same plants as described in Table 1 and 2.

<u>Variety</u>	Flag Leaf		Subtending Leaf		Vegetative Tiller Leaf	
	Length (cm)	Width (mm)	Length (cm)	Width (mm)	Length (cm)	Width (mm)
Bonsai	10.74	4.41	16.86	6.06	30.46	6.28
SR 8600	12.54	5.50	20.91	6.64	39.40	6.60
Crewcut II	12.68	4.68	20.02	5.88	33.53	5.64
Crewcut	13.34	5.79	22.36	6.91	42.23	6.65
SR 8500	13.59	4.89	21.31	6.10	35.24	6.49
Silverado	13.87	5.16	22.16	6.51	40.74	6.42
MiniMustang	13.90	4.56	21.97	6.09	41.34	5.92
Titan Ltd.	14.16	4.84	23.41	5.85	40.53	6.09
Rebel Jr.	14.72	5.99	23.66	7.16	42.45	6.28
Titan	14.88	5.68	25.78	7.18	48.58	6.87
Shortstop	14.98	5.55	22.99	6.66	38.06	6.34
Rendition	15.15	5.42	23.61	5.88	35.82	6.25
Grande	15.20	6.01	22.92	7.28	40.82	6.33
SR 8210	15.56	5.55	27.09	6.94	50.90	6.98
SR 8200	15.78	6.26	26.59	7.55	48.78	6.75
Kentucky 31 NE	16.24	6.19	28.00	8.28	54.70	8.93
Kentucky 31 E	16.29	6.81	28.29	9.07	53.92	8.24
Rebel II	17.3 9	6.13	29.10	7.34	41.21	7.16
Bonanza	17.85	6.61	28.82	8.01	49.21	7.71
LSD@5%	1.86	0.55	2.01	0.54	4.43	0.54

Table 6. Leaf characteristics of tall fescue varieties in 2001 near Corvallis, OR. The plants are those described in Tables 1 and 3. The leaves were measured on the reproductive tillers utilized in Table 4. The vegetative leaf was measured as the last fully expanded leaf on vegetative tillers on the same plants as described in Table 1 and 2. Less varieties and experimentals were measured in 2001.

Variety	Flag Leaf		Subtending Leaf		Vegetative Tiller Leaf	
	Length (cm)	Width (mm)	Length (cm)	Width (mm)	Length (cm)	Width (mm)
Rendition	14.05	3.95	19.56	4.83	26.47	6.36
Bonsai	14.48	3.95	19.23	4.60	30.52	6.46
Titan Ltd.	15.46	4.03	25.32	5.39	32.94	6.18
Silverado	16.59	4.84	24.84	5.23	38.08	6.05
Titan	16.72	5.69	24.77	6.88	43.47	7.83
Crewcut	16.73	4.90	25.54	5 <i>.</i> 71	42.01	6.36
Kentucky 31 E	17.20	6.12	24.74	7.36	49.28	8.10
Shortstop	17.54	5.32	22.75	5.99	40.07	6.41
MiniMustang	17.60	5.05	21.47	5.71	36.62	6.41
SR 8600	17.93	5.62	22.52	5.74	33.85	6.03
Rebel Jr.	18.28	5.58	22.84	5.83	39.89	6.44
LSD@5%	2.30	0.55	2.58	0.59	3.32	0.54

Table . Performance of tall fescue cultivars and selections in a turf trial seeded in August 2000 at Adelphia, NJ.

	Cultivar or Selection	Turf Quality¹ 2001 Avg.	Seedling Height ² Sept. 2000	Establishment ³ Sept. 2000	Leaf Spot⁴ Nov. 2000	Brown Patch ^s 2001 Avg.
·					<u> </u>	
1	DOL comp	6.8	8.3	7.0	6.3	7.2
2	SBM comp	6.5	7.3	7.3	5.7	6.3
3	RB3 comp	6.4	7.3	6.3	6.0	5.7
4	OD3 comp	6.2	6.3	6.3	5.7	7.3
5	10,001 comp	6.2	6.7	6.7	5.7	6.7
6	OD4 comp	6.1	7.7	6.3	5.0	7.2
7	P58	6.1	6.7	7.3	7.3	6.5
8	TF-33	6.1	7.3	6.3	5.7	5.7
9	TF-34	6.1	7.3	6.7	6.3	4.5
10	2nd Millennium	6.1	7.0	6.0	5.3	7.2
11	Syn 578	6.1	5.3	6.3	5.3	7.0
12	Forte	6.1	6.7	6.3	6.0	6.8
13	TF-35	6.1 ·	7.7	6.0	6.0	6.8
14	Justice	6.0	7.0	6.3	4.0	7.5
15	EA 171	6.0	7.7	5.7	4.3	5.3
16	OD1 comp	6.0	6.7	6.7	5.3	6.7
17	OD2 comp	5.9	6.0	6.0	4.7	7.3
18	00 GFA	5.9	5.7	6.0	5.3	6.5
19	Syn 5K1	5.9	6.3	6.0	6.0	6.3
20	SR 8600	5.9	5.0	6.7	6.3	5.7
21	Bingo	5.9	6.7	6.7	4.7	5.0
22	Biltmore	5.8	6.7	5.3	5.3	5.7
23	Syn 5T2	5.8	6.3	6.0	4.7	5.7
24	Syn 5BAB	5.7	6.3	6.3	5.3	6.8
25	00-BFA	5.7	6.0	7.0	4.7	5.2
	Matador	5.6	8.0	5.7	6.3	4.7
	Syn 5KU	5.6	₹7.7	6.3	5.3	6.2
	BE 4	5.6	8.0	6.3	4.7	5.7
	Mustang III	5.6	6.0	6.3	5.0	5.7
30	Rendition	5.6	6.7	6.7	5.7	5.8
	00-H FA	5.6	6.3	6.0	5.0	5.8
	Syn 5DWF	5.6	7.7	6.7	6.0	5.7
	E-97	5.6	8.3	6.7	4.0	5.0
	FA6-91	5.5	6.7	5.7	5.0	5.8
35	Syn-R54M-00	5.5	8.3	6.3	5.3	3.3

(Continued)

200300174

Table $\rlap{/}{n}$ (continued).

	Cultivar or Selection	Turf Quality¹ 2001 Avg.	Seedling Height ² Sept. 2000	Establishment ³ Sept. 2000	Leaf Spot⁴ Nov. 2000	Brown Patch⁵ 2001 Avg.
36	Syn 5NAS	5.5	6.7	6.3	5.7	5.3
37	SRX 8DDMPP	5.4	6.7	5.7	4.7	6.3
38	MA 176	5.4	7.3	5.7	3.7	6.2
39	Syn 5BEH	5.4	6.3	5.7	5.3	6.0
40	Syn 5BZ	5.4	6.0	5.3	5.0	6.0
41	Southern Comfort	5.4	6.7	6.0	5.0	5.0
42	DLSD	5.4	6.0	6.0	6.0	5.0
43	Syn 5MP	5.4	7.0	6.7	5.7	4.7
44	57E	5.4	6.3	5.7	6.3	4.8
45	TF H-97	5.4	7.0	6.3	5.3	4.2
46	00-J FA	5.4	7.3	6.7	6.3	4.8
47	Picasso	5.4	7.0	6.0	5.3	3.8
48	EA 180	5.3	7.7	6.0	3.3	5.5
49	TF-34	5.3	8.0	6.3	5.0	4.8
50	Syn 5H2	5.3	5.7	5.7	4.7	6.2
51	SRX 8 FFT	5.3	6.7	6.0	4.7	5.5
52	00-CFA	5.3	5.3	5.7	3.7	5.2
53	5BE	5.3	6.3	6.7	4.7	5.5
54	00-A FA	5.3	6.3	6.7	4.7	3.8
55	SRX 8601 E	5.2	5.7	6.0	5.3	6.3
56	Syn 5A3	5.2	6.7	5.7	5.3	5.7
57	Rembrandt	5.2	6.0	6.0	4.7	5.2
58	SRX 8BPDDE	5.2	6.0	6.3	4.7	4.7
59	5301	5.2	.•	•	•	6.5
60	Santa Fe	5.2	7.0	5.7	2.0	5.0
61	Crewcut II	5.2	6.0	5.7	5.3	3.7
62	BE1	5.1	_. 6.3	5.7	5.0	6.7
63	Plantation	5.1	⁷ 5.7	6.7	4.7	5.0
64	MA 157	5.1	7.3	5.7	4.0	4.8
65	TF J-97	5.1	7.0	6.0	4.7	4.5
66	EA 172	5.1	6.3	5.7	3.0	4.2
67	Laramie	5.1	6.0	6.0	5.3	3.7
68	Sun Pro	5.1	6.3	5.3	5.3	5.0
69	MA 160	5.1	7.7	5.0	4.0	4.3
70	Syn-R5JM-00	5.1	7.7	6.3	5.0	2.8

	Cultivar or Selection	Turf Quality¹ 2001 Avg.	Seedling Height ² Sept. 2000	Establishment ³ Sept. 2000	Leaf Spot⁴ Nov. 2000	Brown Patch ^s 2001 Avg.
71	CAE comp	5.1	5.7	5.7	4.7	6.5
72	Syn TUO	5.1	6.7	6.7	4.0	5.2
	Bravo	5.1	4.3	6.3	5.0	4.2
	EA 155	5.1	6.7	5.7	4.0	4.2
75	MC1	5.0	6.0	5.7	5.0	7.0
	Syn 5S2	5.0	5.7	5.7	5.7	5.5
	TF-41	5.0	7.3	6.7	5.0	5.7
	Syn 5CH	4.9	6.0	5.7	5.0	6.0
	Pick FA B93	4.9	6.7	5.3	5.0	5.5
80	Syn BRO	4.9	5.3	6.3	4.3	5.3
	SRX 8DDEOO	· 4.9	6.3	5.7	4.3	4.2
	8 S M2	4.9	6.0	5.7	3.3	5.2
	MA 127 Forbes	4.9	6.7	5.7	2.7	5.2
	P89 * SpL	4.9	5.3	6.3	5.0	5.2
85	SRX 8 BPDDNE	4.9	5.3	6.3	2.7	5.0
	ORE-00TF	4.9	4.7	6.0	4.3	4.3
	Millennium	4.8	5.7	6.3	4.0	5.3
	EA 163	4.8	6.7	5.3	3.3	4.7
	Pure Gold	4.8	7.0	6.7	4.3	3.5
90	Syn 5G9	4.8	6.3	5.3	6.0	5.7
	00-I FA	4.8	6.7	6.0	4.0	4.8
	00-D FA	- 4.8	5.0	5.7	4.0	6.3
	MA 177	4.8	7.7	5.3	3.3	4.7
	SRX 8EDFF Shortstop II	4.7 4.7	4.7 7.0	5.3 5.3	4.7 5.3	4.5
	•			5.3	ე. ა	4.2
	ΓF-40	4.7	5.7	6.0	3.3	4.2
	SRX 8CDEW	4.7	7.0	6.7	3.3	5.0
	SRX 8MO961	4.7	5.7	6.3	5.0	4.5
	Prospect	4.6	5.7	6.0	4.3	6.5
1 00	MA 165	4.6	7.0	5.0	4.0	3.5
	Apache II	4.6	5.7	6.0	5.0	5.0
	Coronado Gold	4.6	4.7	6.0	4.7	4.8
	RT-95	4.6	5.3	6.0	3.3	4.5
	SR 8500	4.6	5.7	6.0	5.0	5.2
105	SS bulk M2	4.6	6.3	6.3	4.7	3.7

Table 🖡	(continued).		•		: ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° ° °	
	Cultivar or Selection	Turf Quality ^t 2001 Avg.	Seedling Height ² Sept. 2000	Establishment ³ Sept. 2000	Leaf Spot ⁴ Nov. 2000	Brown Patch ⁵ 2001 Avg.
100	MA 170	4.6	6.3	5.3	3.0	3.3
106 107	MA 178 MA 138 JSC	4.5	6.0	6.3	3.7	5.0
107	80P22	4.5 4.5	7.3	5.7	3.0	4.3
109	8RF2	4.5	6.3	6.0	3.0	4.8
110	GS bulk E1	4.5	4.7	6.0	4.0	4.5
111	Coronado	4.5	6.3	6.3	4.7	2.8
112	00-E FA	4.5	5.3	5.3	3.0	5.2
113	Houndog 5	4.5	4.0	5.7	4.0	4.3
114	Olympic Gold	4.5	4.7	6.0	4.7	4.2
115	MA 158	4.4	8.0	4.7	2.7	5.7
116	Rebel Jr.	4.4	3.7	5.7	3.3	4.7
117	Tarheel	4.4	5.0	5.7	3.7	3.5
118	5UD	4.4	6.3	5.7	4.7	4.7
119	Lancer	4.3	5.0	5.3	4.3	4.3
120	Tomahawk E+	4.3	4.0	6.0	4.7	3.8
121	MA 98 Mtn. View	4.3	6.3	6.0	2.0	4.0
122	TF-43	4.2	5.7	5.3	3.0	5.2
123	P89 * SpE	4.2	6.3	6.0	4.3	3.7
124	TF-42	4.2	5.3	5.3	2.3	5.8
125	Talisman	4.2	6.3	6.0	4.7	4.5
126	Syn 5HUO	4.1	5.7	6.7	3.3	5.7
127	Tomahawk	4.1	3.3	5.7	2.7	4.7
128	Crossfire II	4.1	5.7	5.7	4.3	4.5
129	T991-00	4.0	6.7	5.3	4.3	3.5 5.2
130	Regiment	4.0	2.7	5.0	4.0	5.2
131	SRX 8 MO94	4.0	4.3	6.3	3.3	4.8 = 0
132	Grande	3.9	3.3	6.3	3.0	5.8
133	Crewcut	3.9	<i>,</i> 5.0	5.7	4.7	4.2
134	D5 ATF 00-6	3.9			•	6.7
135	SRX LJHH	3.8	5.0	5.7	3.7	5.3
136	Bonanza II	3.8	3.7	5.3	2.7	3.7
137 -		3.8	4.3	6.0	3.3	3.7
138	Wolfpack	3.7	4.7	4.7	3.7	5.7 5.2
139	Mustang II	3.7	2.7	5.3	2.7	5.3
140	Hilitop TF	3.6	2.3	5.0	2.3	4.0

(Continued)

Table # (continued).

	Cultivar or Selection	Turf Quality¹ 2001 Avg.	Seedling Height ² Sept. 2000	Establishment³ Sept. 2000	Leaf Spot ⁴ Nov. 2000	Brown Patch ^s 2001 Avg.
1	Eldorado	3.6	2.7	4.3	3.7	5.5
?	Confederate	3.6	5.0	4.3	4.0	5.0
3	GT 2K	3.0	4.3	5.0	1.3	4.5
1	Kentucky 31	1.3	1.0	4.0	1.7	5.2
5	Torpedo	1.0	1.0	3.3	1.0	5.7
	LSD at 5% =	0.7	1.2	1.1	1.2	2.1

^{&#}x27;9 = best turf quality

²9 = shortest seedling height

³9 = best establishment

^{49 =} least leaf spot

⁵9 = least brown patch (average of two ratings taken 8-13-01 and 8-17-01)

REPRODUCE LOCALLY. Include form number and edition date on all	reproductions. F	ORM APPROVED - OMB No. 0581-0055		
U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE EXHIBIT E	Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). The information is held confidential until the certificate is issued (7 U.S.C. 2426).			
STATEMENT OF THE BASIS OF OWNERSHIP	O TEMPODADY DESCRIPTION	LO MADIETY MANE		
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME		
Seed Research of Oregon	SRX 8V9	Rendition		
4. ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP, and Country) 27630 Llewellyn Rd.	5. TELEPHONE (Include area code)	6. FAX (Include area code)		
Corvallis, OR 97333	541-757-2663 7. PVPO NUMBER	541-752-2065		
		00174		
8. Does the applicant own all rights to the variety? Mark an "X" in the	e anoropriate block if no please expla	in. YES NO		
A royalty is split between SRO a 9. Is the applicant (individual or company) a U.S. national or a U.S. b	nd Rutgers Univers	ity		
. ,				
10. Is the applicant the original owner?	NO If no, please answer one	of the following:		
ا الما				
a. If the original rights to variety were owned by individual(s), is (are) the original owner(s) a U.S. National NO If no, give name of countr			
b. If the original rights to variety were owned by a company(ies), YES [, is (are) the original owner(s) a U.S. bar			
11. Additional explanation on ownership (Trace ownership from origin Germplasm acquired from Rutgers	nal breeder to current owner. Use the re University underwe	overse for extra space if needed): nt cycles of selection		
by Dr. Leah A. Brilman of Seed R	esearch of Oregon.			
PLEASE NOTE:				
Plant variety protection can only be afforded to the owners (not licens	sees) who meet the following criteria:			
If the rights to the variety are owned by the original breeder, that penational of a country which affords similar protection to nationals of a country which affords similar protection.				
If the rights to the variety are owned by the company which employ nationals of a UPOV member country, or owned by nationals of a o genus and species.				
3. If the applicant is an owner who is not the original owner, both the	original owner and the applicant must m	eet one of the above criteria.		
The original breeder/owner may be the individual or company who did Act for definitions.	rected the final breeding. See Section 4	1(a)(2) of the Plant Variety Protection		
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, control number. The valid OMB control number for this information collection is 0581-0055, including the time for reviewing the instructions, searching existing data sources, gathering a	The time required to complete this information collect	tion is estimated to average 0.1 hour per response,		
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and a marital or family status, political beliefs, parental status, or protected genetic information. (N. companying the property information, Paralle, Agraille, Lange print, supplies on a context U.	ctivities on the basis of race, color, national origin, ge ot all prohibited bases apply to all programs.) Person	ander, religion, age, disability, sexual orientation, ns with disabilities who require alternative means for		

To file a complaint of discrimination, write USDA, Director, Office of Civil Rights, Room 326-W, Whitten Building, 14th and Independence Avenue, SW, Washington, D.C. 20250-9410 or call (202) 720-5964 (voice and TDD). USDA is an equal opportunity provide and employer.